

# AMBUSH

AMBUSH is an emulsifiable concentrate formulation containing 100 g/l (11% w/w) cypermethrin

AMBUSH is a broad spectrum contact insecticide for the control of agricultural and horticultural insect pests.

## RISK AND SAFETY INFORMATION

**IRRITATING** TO THE RESPIRATORY SYSTEM.

**HARMFUL** IF SWALLOWED, MAY CAUSE LUNG DAMAGE.

**VERY TOXIC** TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT. **FLAMMABLE.**

REPEATED EXPOSE MAY CAUSE SKIN DRYNESS OR CRACKING. VAPOURS MAY CAUSE DROWSINESS AND DIZZINESS.



PCS No. 04509

Contents: 1LE

Batch No: see packaging

**Registration holder:**  
DHM Agrochemicals  
Castle Lodge  
Kilgobbin Road  
Dublin 18

Tel: 01 2952377 - Fax: 01 2959399

**Marketed by:**  
DHM Agrochemicals  
Castle Lodge  
Kilgobbin Road  
Dublin 18

Tel: 01 2952377 - Fax: 01 2959399

## COMPLIANCE WITH THE FOLLOWING CONDITIONS OF USE IS A LEGAL REQUIREMENT FOR USE ONLY AS AN AGRICULTURAL / HORTICULTURAL INSECTICIDE

Crop: Pests	Rate (ml/ha)	Remarks/Timing
<u>Winter and spring cereals</u> (wheat, oats, barley): Barley Yellow Dwarf Virus, Aphids	250	Apply October–November after crop emergence.
<u>Root crops (sugar beet, fodder beet, red beet, mangels):</u> Cut worms	250	Apply at egg hatching, repeat 14 days later.
<u>Brassicac:</u> Caterpillars	250	Spray at the first sign of attack.
<u>Field beans:</u> Pea and Bean Weevil	250	Spray at first sign of leaf notching in April/May when adults are active.
<u>Oilseed Rape:</u> Cabbage stem flea beetle	250	Adults, apply at first sign of damage usually in September. To control further larval attack, apply late November.
Rape Winter Stem Weevil	250	Apply in October or when adults are seen in crop.
<u>Tomato &amp; Pepper (Glasshouse):</u> White fly and leaf miner	500-750	Apply when pests first seen. Repeat at 4-5 day interval until pest disappears.
<u>Pears:</u> pear sucker	350	Apply February/March to kill adults before egg laying. Repeat in summer if necessary.
<u>Apples pre-blossom:</u> winter moth, Tortrix, Caterpillar, aphids, suckers	280	Apply at green cluster to pink bud.
<u>Apples post-blossom:</u> sawfly, capsid	280	Petal fall-repeat if re-infestation occurs.
Fruitlet Mining Tortrix	280	Apply 14 days after petal fall.
Tortrix and codling moths	280	Apply in June/July or as per indications by moth trap catches.

READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE. TO AVOID RISKS TO MAN AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE.

## DIRECTIONS FOR USE

Add concentrate to water as required and agitate. For high volume recommendations on fruit, apply just short of run off (approximately 1000 l/ha). For root crops and Brassicas, apply 600 l water/ha. For other crops, apply a low volume of water (approximately 200 l/ha). Ambush is not phytotoxic when used as recommended and is compatible with a wide range of products. Ambush does not taint brassicas and peas when applied up to ten days before harvesting. However, the processors should be consulted before application. Ambush acts by contact, therefore ensure thorough spray cover. Always avoid spraying under stress conditions.

## SAFETY PRECAUTIONS

### Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES (NEOPRENE OR NITRILE) AND FACE PROTECTION (FACESHIELD) when opening the container, transferring the contents from one container to another, diluting, mixing, filling chemical tank, adjusting apparatus after filling, washing out containers after use.

<sup>1</sup>at least 0.5mm thickness

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

TAKE OFF IMMEDIATELY all contaminated clothing.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

### Environmental protection

**EXTREMELY DANGEROUS TO FISH OR OTHER AQUATIC LIFE:** Do not contaminate surface waters or ditches with chemical or used container.

Dangerous to bees.

### Storage and disposal

DO NOT RE-USE CONTAINER for any purpose.

PROTECT FROM FROST.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

### MEDICAL ADVICE

#### First Aid

In case of skin contact remove all contaminated clothing and wash the patient thoroughly with plenty of water and soap. In case of eye contact, flush eyes with copious amounts of water for at least 15 minutes and seek medical advice. If by inhalation move to fresh air, administer oxygen if necessary. Keep the patient warm and quiet. If swallowed, do not induce vomiting; seek medical advice immediately and show this container or label. Gastric lavage should be carried out with protection of the airways. Keep the patient at absolute rest and seek medical assistance immediately. Do not induce vomiting to an unconscious patient.

Note to doctor: No specific antidote known. Symptomatic treatment.

### CONDITIONS OF SALE

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.