

DimethoX

Contains 400 g/litre (38% w/w) dimethoate as an emulsifiable concentrate

A contact and systemic insecticide for the control of aphids and certain other pests
on wheat, potatoes and seed crops of sugar beet and mangels

FOR PROFESSIONAL USE ONLY

Product identifier: Dimethoate 400 g/l EC

Contains dimethoate, cyclohexanone, hydrocarbons, C9, aromatics and maleic anhydride



Danger

Flammable liquid and vapour.

Harmful if swallowed.

May be fatal if swallowed and enters airways.

May cause an allergic skin reaction.

Harmful if inhaled.

Very toxic to aquatic life with long lasting effects.

Avoid breathing vapours.

Wear protective gloves and eye/face protection.

IF SWALLOWED: Rinse mouth.

IF ON SKIN (or hair): Remove immediately all contaminated clothing. Wash with plenty of soap and water.

Immediately call a POISON CENTER or doctor/physician.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads

To avoid risks to human health and the environment, comply with the instructions of use.

PCS No 05118

CONTENTS: 5 LITRES e

(Packed in Cartons of 4 x 5L)



APPROVAL HOLDER AND MARKETING COMPANY: DHM AGROCHEMICALS LTD.,
CASTLE LODGE, KILGOBBIN ROAD - DUBLIN 18. TEL: 01/2952377 - FAX : 01/2959399

PROTECT FROM FROST

6/22504-FL

DimethoX

DIMETHOX DIRECTIONS FOR USE

CROP	PESTS	WHEN TO APPLY	RATE
Potato Seed crops,	Aphid Excluding <i>Myzus persicae</i>	At 75% emergence. Repeat every 14 days to burning off.	840 ml in 200-1000 litres/ha water.
Wheat	Aphid	Spray when aphids are first seen. Do not apply later than 21 days before harvest	840 ml in 200-1000 litres/ha water.
	Wheat bulb fly	Spray as soon as damage is first seen usually early March	1680 ml in 200-1000 litres/ha water.
<u>Stecklings</u>			
Seed crops of: Sugar beet Mangels	Black aphid (excludes <i>Myzus persicae</i>)	When pest first seen. If stecklings are grown under a cover crop, spray once the cover crop is harvested. Repeat every 14 days to the end of October.	840 ml in 200-1000 litres/ha water.
<u>Harvest year</u>			
		When pest first seen.	1050 ml in 400-1000 litres/ha water.

Mixing

Half fill the spray tank with water. Add the required amount of insecticide, top up the tank with water and agitate.

Consult Processor

Before using on crops for processing.

Insect Resistance

Some insects, particularly aphids, have in some areas become resistant to dimethoate and other organophosphorus insecticides. In such circumstances, good control from Dimethoate will not be obtained.

Storage

Store in a cool dry place, away from foodstuffs, seeds and other chemicals. Flammable. Protect from frost.

Do not contaminate water with the product or its container.

(Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads)

READ CAREFULLY THE RECOMMENDATIONS FOR USE PROVIDED WITH THIS CONTAINER.