

Headland Relay

Contains 200 g/l (16.5% w/w) mecoprop-p, 200 g/l (16.5% w/w) MCPA and 25 g/l (2.0% w/w) dicamba
A selective weedkiller for use in grass leys, permanent pastures and amenity turf to control a broad spectrum of annual and perennial weeds



RISK AND SAFETY PHRASES

Harmful if swallowed.

Causes serious eye damage.

Toxic to aquatic life with long lasting effects.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

Avoid release to the environment.

IF SWALLOWED: Call a POISON CENTER or doctor/ physician if you feel unwell.

Rinse mouth.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Dispose of contents / container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which may be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To avoid risks to human health and the environment, comply with the instructions for use.

Crop	Maximum individual dose	Maximum Number of treatments	Latest time of application
Permanent Pasture Grass leys	7.0 l/ha	2 per year	7 days before cutting or grazing
Amenity grassland	5.0 l/ha	2 per year	-

Registration Number PCS 90728

CONTENTS: 5 LITRES

 **Headland**

Batch No: see neck of bottle

 **CHEMINOVA**



Headland Agrochemicals Ltd., Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.
 Telephone: 01244 537370 Fax: 01244 532097 E-mail: enquiry@headlandgroup.com www.headland-ag.co.uk

Headland Relay

FOR USE ONLY AS AN AGRICULTURAL / AMENITY HERBICIDE

RATE OF USE IN GRASS LEYS AND PERMANENT PASTURE

Use 5 litres/ha (3½ pints/acre) in 200 litres of water/ha (18 gallons/acre).

For improved control of Docks, Ragwort and other weeds in grass leys and permanent pastures where the grass has been established for at least 18 months, use 7 litres/ha (5 pints/acre) in at least 200 litres of water /ha (18 gallons/acre). Note: Do not use this higher rate in amenity turf.

TIMING IN GRASS LEYS AND PERMANENT PASTURE

Apply from spring to early October when weeds are growing actively, preferably at the rosette stage. Larger weeds, particularly Docks, Ragwort and Soft Rush may need a further application. In cold weather or in slow growing conditions the inclusion of Headland Intake penetrant will be of benefit.

RATE OF USE IN AMENITY TURF

Use 5 litres/ha (3½ pints/acre) in 200 litres of water/ha (18 gallons/acre).

TIMING IN AMENITY TURF

Apply at any time between April and early October when both grass and weeds are growing strongly. Best results will be obtained in fine, warm weather when the soil is moist.

If possible, apply a nitrogen fertiliser 7-14 days before treatment.

Do not use on new grass for 6 months after establishment.

MIXING AND APPLICATION

Make sure the sprayer has been properly cleaned from any previous use. Half-fill the spray tank with clean water and add the required amount of Headland Relay whilst agitating. Then add the remaining quantity of water and apply without delay. If the machine is stopped for any time, agitate thoroughly before restarting spraying.

After use wash out the sprayer thoroughly using a suitable spray tank decontaminant.

COMPATIBILITY

Headland Relay is compatible with many commonly used pesticides and micronutrients. Your Headland distributor will be able to answer any specific questions about compatibility. DO NOT MIX WITH MANGANESE SULPHATE OR FERROUS SULPHATE.

CAUTION

AVOID SPRAYING when wind could cause drift.

AVOID DRIFT ONTO SENSITIVE CROPS such as beet, brassicas, lettuce, tomatoes, peas, potatoes, fruit or ornamentals.

DO NOT SPRAY where clovers or other legumes are an important part of the sward.

DO NOT ROLL OR HARROW CROPS within 7 days before or after spraying

DO NOT CUT OR GRAZE grass leys or permanent pasture for at least 7 days after spraying, or 14 days if poisonous weeds such as Ragwort are present. Dead foliage of poisonous weeds must have dried out and become unpalatable. In fields intended for hay or silage, delay spraying until after crops have been taken.

DO NOT CLOSE MOW amenity grass 1 day before or 4 days after spraying. On very fine closely mown turf this interval may be reduced to 1 day.

DO NOT USE grass cuttings for mulching for at least 6 months after application. Cuttings may be composted and used not less than 6 months later.

DO NOT SPRAY when rain is imminent. In changeable weather, tank-mixing with Headland Guard will improve rain-fastness.

PROTECT FROM FROST

Made in the EU

PCS No. 90728

Headland Agrochemicals Ltd., Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.
Telephone: 01244 537370 Fax: 01244 532097 E-mail: enquiry@headlandgroup.com www.headland-ag.co.uk