



## READ LABEL CAREFULLY BEFORE USE

### IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL/HORTICULTURALMOLLUSCID				
Crop	Max. individual dose	Max. no. of applications	Max.total dose	Latest timing of application
Wheat, barley, oats, rye & triticale	4 kg/ha		12 kg/ha	GS 29 (end of tillering)
Oilseed rape	4 kg/ha		12 kg/ha	GS 29 (9 or more side buds visible)
Maize	4 kg/ha		12 kg/ha	GS 15 (5th leaf unfolded)
Lettuce	4 kg/ha		12 kg/ha	GS 39 (rosettes complete)
Field Berries (Blackberry, Blackcurrant, Blueberry, Cranberry, Gooseberry, Loganberry, Raspberry, Redcurrant, Whitecurrant)	4 kg/ha		12 kg/ha	GS 69 (End of flowering)
Pome fruit & Stone fruit (apple, pear, plum, cherry)	4 kg/ha		12 kg/ha	GS 69 (End of flowering)
Grapes (Wine and Table)	4 kg/ha		12 kg/ha	GS 69 (End of flowering)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

#### Safety Information

If medical advice is needed, have product container or label at hand. Keep out of the reach of children.

Do not get in eyes, on skin, or on clothing.

Wear protective gloves.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use**

PCS No. 04960

#### PRECAUTIONS

If you feel unwell seek medical advice immediately (show the label where possible).

In case of accident, call the National Poisons Information Centre: Tel: +353 (0)1 809 2566 or +353 (0)1 837 9964

Do not contaminate water with the product or its container. Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms do not apply on impermeable surfaces such as asphalt, concrete, cobblestones, railway tracks and other situations with a high risk of run-off.

To protect wild birds and wild mammals remove spillages.

Keep poultry out of treated areas for at least 7 days following application.

Protect from frost, store in a dry, cool, frost-free ventilated area.

24 hour emergency number – National Poisons Centre: +353 (01) 809 2566 or +353 (01) 837 9964

Net content:

Batch number: see container

Registration no.:

**Registration holder:** Sharda Europe b.v.b.a, Jozef Mertensstraat 142, 1702 Dilbeek, Belgium

**Distributed by:**

**Phone no: +91 22 6678 2800**

**Emergency number: +91 22 6678 2800 (8.00 – 16.00h)**

Registered logos of Sharda Cropchem Ltd.



**Sharda Cropchem**

SHARDA CROPCHEM LIMITED  
Dornic Holm 29<sup>th</sup> Road, Bandra (West)  
Mumbai - 400050, India, Tel. +39 02 66101029

TRAXX is a molluscicide for the control of slugs in crops.

#### DIRECTIONS FOR USE:

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

#### RESTRICTIONS

Professional use only

DO NOT apply before heavy rainfall, which may reduce the persistence of granules. Late application of pellets to broad-leaved plants may result in lodging of the pellets in foliage. Care should be taken to avoid this during application in edible crops.

Keep dogs out of the field during application. When the applicators stops/starts, ensure no small heaps of pellets are left on the soil surface as they may be consumed by dogs and other animals. Spillages must be buried or removed.

#### APPLICATION RECOMMENDATIONS

Best results are achieved when application is made during mild, damp weather when slugs are the most active. When slug attack is heavy or prolonged, repeat treatments may be required.

#### All crops

4 kg/ha broadcast evenly over soil or standing crop using any suitable slug pellet applicator. For crops to be drilled best results are from application a few days before drilling. Alternatively, apply just before or just after drilling. For emerged or established crops, apply immediately damage appears. Do not apply more than three times per season (maximum of 12 kg/ha) Minimum interval between applications 7days

#### COMPANY ADVISORY INFORMATION

Slug pellets are pesticides, keep pesticides out of water

- Pellets should not fall within 5m of the top of the bank of a watercourse or ditch. To achieve this, users should determine the spread width of the applicator with the product to be applied prior to any application (to the rear as well as the side of the applicator).
- o Calibrate your applicator before use
- o Fill and clean application equipment in the field
- o Do not fill or clean down in the yard
- o Avoid applying slug pellets when heavy rain is forecast
- o Check that product packaging is intact and clear up spillages immediately