

**PROTECT FROM FROST
SHAKE WELL BEFORE USE**

WINGER provides moderate control of powdery mildew on winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

CONDITIONS FOR USE

FOR USE AS AN AGRICULTURAL FUNGICIDE

CROP	Max single dose (litres/hectare/crop)	Max no of applications	Max total dose (litres/hectare/crop)	Latest timing of application
Winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.	0.5l/ha	-	1.0	Before beginning of flowering: first anthers visible (GS 61)

Additional Safety Information

Operator protection

Wash splashes from skin and eyes immediately.
Wash hands and exposed skin before meals and work.
Wear suitable protective clothing, gloves and eye/face protection.
Harmful to livestock. Keep livestock out of treated areas for at least 7 days.
In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.
Do not breathe spray.

Storage and disposal

Keep in original container, tightly closed in a safe place.
Wash out containers thoroughly, preferably using an integrated pressure rinsing device, or manually rinse three times. Add washings to the sprayer at the time of filling. Complete filling to the required volume and continue to agitate throughout the spraying operation.

Restrictions

Do not apply more than 3 times to any one crop.
WINGER should only be used up to the full ear emergence (GS 59) stage on all crops.
Disease control may be reduced if strains of the pathogen less sensitive to WINGER develop.

WINGER

WINGER is an emulsifiable concentrate containing 750 grams fenpropidin (81.8% w/w) per litre.

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

WINGER provides moderate control of powdery mildew on winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.



FOR PROFESSIONAL USE ONLY

To avoid risks to human health and the environment, comply with the instructions for use.
WINGER is an emulsifiable concentrate containing 750 g/l (81.8% w/w) fenpropidin.

Warning.
Harmful if swallowed.
Causes serious eye irritation.
Harmful if inhaled.
May cause respiratory irritation.
Very toxic to aquatic life with long lasting effects.

Keep out of reach of children.
Avoid breathing dust/fume/gas/mist/vapours/spray.
Wear protective gloves/protective clothing/eye protection/face protection.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Call a POISON CENTRE or doctor/physician if you feel unwell.
If eye irritation persists: Get medical advice/attention.
Collect spillage.
Store in a well-ventilated place. Keep container tightly closed.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.
To protect aquatic organisms respect an unsprayed buffer zone of 15m to surface water bodies.
Contains fenpropidin. May produce an allergic reaction.

PCS No: 05807



Authorisation Holder	Marketing Company	Net contents: 1 litre Product code / Print date / Batch Number
DHM Agrochemicals Ltd Castle Lodge Kilgobbin Road Dublin 18 Tel: 01 2952377	DHM Agrochemicals Ltd Castle Lodge Kilgobbin Road Dublin 18 Tel: 01 2952377	

DIRECTIONS FOR USE

IMPORTANT: this information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

CROPS

WINGER may be used on all varieties of winter wheat, spring wheat, winter barley, spring barley, oats, rye and triticale.

DISEASE CONTROL

For moderate control of powdery mildew apply during the early stages of disease development. Best results will be achieved when not more than 5% of the third leaf from the top of the plant is infected. A repeat spray may be necessary during prolonged or severe attacks.

APPLICATION RATE

Apply WINGER at 0.5 litres per hectare.

MIXING AND SPRAYING

Make sure the sprayer is set to give an even application at the correct volume.
Fill the sprayer tank with half the required quantity of clean water and begin agitation. Add the required quantity of WINGER and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.
Thoroughly wash all spray equipment with water immediately after use.

APPLICATION

Volume

Apply WINGER in a recommended 200 litres of water per hectare.

COMPATIBILITY

The following products may be tank-mixed with WINGER provided the timing is correct for both WINGER and the partner in the mixture.
The products should be added separately to the bulk of the water in the spray tank. Continuous agitation should be maintained and the products used immediately after mixing.

Fungicides

CT 500

Herbicides

Approved salt formulations of:
MCPA
Mecoprop

Growth Regulators

Approved formulations containing chlormequat or chlormequat and choline chloride such as K2, Ceraide and Manipulator.

For further information or compatibility, contact DHM Agrochemicals Ltd.

ULBDHWING1L (03-17)