

An emulsifiable concentrate formulation containing 637 g/L of methylated rape seed oil. FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL/FORESTRY WETTING AND SPREADING AGENT

STATUTORY CONDITIONS RELATING TO USE

Follow the statutory conditions of use of the pesticide

An emulsifiable spray adjuvant specially formulated to enhance the wetting and spreading of herbicides, fungicides and crop nutrition products. UFI No: 0AU0-F104-500T-XHK8

Maximum Concentration:

All crops except fodder beet 0.5% of spray solution Fodder beet: 0.85% of spray solution

Maximum Number of treatments: Follow the statutory conditions of use of the pesticide

Latest timings:

1. On cereals with morpholine or triazole fungicides:

On crops not intended for human or animal consumption: Follow the statutory conditions of use of the pesticide

With all approved pesticides applied to edible crops at Follow the statutory conditions of use of the pesticide

half or less than half rate

With approved pesticides on other crops up to and Including the growth stages given below

CROP	LATEST TIME OF APPLICATION
Apples/pears Strawberries Raspberries	Fruit 5 - 10 mm First fruit development First fruit set
Blackcurrants	First visible green fruitlet
Cucumber Tomato	First fruit set on first truss
Broad bean, dwarf French bean, field bean, runner bean Peas (combining & vining) Edible podded peas Oilseed rape Linseed Sugar beet Cereals Fonge maize Brassels sprouts Potatoes	First pod set Flat pod (non-edible pods only) Pod set 10% potential pods 10% capsules formed 6 fully expanded leaves 114 inflorescence (GSS2) Tip of tassel visible Lateral buds beginning to develop Tuber initiation
Cauliflower Apricot, plum, peach Hops	Heads beginning to form (width up to 1 cm) First fruit set Cone set

WARNING - SAFETY PRECAUTIONS

Avoid release to the environment.

Wear protective gloves/protective clothing/eye protection/face protection.

IF ON SKIN: Wash with plenty of soap

and water.
If skin irritation or rash occurs:

Get medical advice/attention.

Wash contaminated clothing before reuse.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of

To avoid risk to human health and the environment comply with the instructions for use

OPERATOR PROTECTION

Follow all operator protection instructions for the partner pesticide. Where the personal protective equipment specified on this label is different to that specified on the pesticide label the highest level of protective equipment must be worn.

OTHER SPECIFIC RESTRICTION

Do not use RAPPER with any pesticide which is to be applied in or near water.

PROTECT FROM FROST

CONTENTS: 5 litres PCS No: 92491



DIRECTIONS FOR USE

RAPPER may be used with approved pesticides in accordance with the Conditions of Use. The Conditions of Use of the partner pesticide must also be complied with.

Use RAPPER at a concentration of 0.25 - 0.5% of total spray volume (250 - 500 ml per 100 litres of water or pro rata). When used with beet herbicides, this rate may be increased to 0.85% of water volume if required.

Use the higher rate under difficult conditions such as high weed populations (herbicides), high disease pressures (fungicides), dense canopies, weather stress etc. Do not contaminate water with the productor its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and road

MIXING INSTRUCTIONS

Charge the sprayer with half the recommended volume of water and the pesticide in accordance with the manufacturer's instructions. Add the required amount of RAPPER followed by the remaining water requirement as the last operation. Maintain agitation whilst mixing, travelling and spraying. Clean all equipment thoroughly after use. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste

COMPATIBILITY

RAPPER is compatible with most foliar-applied pesticides. Please consult your distributor for information on specific tank mixes.

STORACE

Protect from frost. Store above 5°C.

Keep in cool, well ventilated suitable pesticide store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs. If RAPPER is subjected to temperatures below 0° C it may thicken and become cloudy. It will return to its normal clear fluid state above 0° C.

OPERATOR PROTECTION

Follow all operator protection instructions for the partner pesticide. Where the personal protective equipment specified on this label is different to that specified on the pesticide label the highest level of protective equipment must be worn. In the event of an emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964

OTHER SPECIFIC RESTRICTION

Do not use RAPPER with any pesticide which is to be applied in or near water.

Rapper is a registered trademark of DHM Agrochemicals Ltd. Castle Lodge, Kilgobbin Rd, Dublin 18. PH: 01 2952377